


ONE STEP PROCESS FOR THE PREPARATION OF ANTICONVULSANT
DERIVATIVES

Cross Reference to Related Application

5 This application claims the benefit of U. S. Provisional Application 60/451,863, filed on March 4, 2003, which is incorporated by reference herein in its entirety.

Field of the Invention


10 The present invention relates to a one step process for the preparation of compounds of the formula (I)



15 wherein X, R¹, R³, R⁴, R⁵ and R⁶ are as hereinafter defined. The compounds of formula (I) are useful for the treatment of epilepsy.

Background of the Invention


The compounds of formula (I)



20 wherein R¹ is selected from the group consisting of hydrogen and C₁₋₄alkyl and wherein X, R³, R⁴, R⁵ and R⁶ are as hereinafter defined, are known compounds that have been found to exhibit anticonvulsant activity and are therefore
25 useful in the treatment of conditions such as epilepsy. These compounds are disclosed in U.S. Patent No. 4,582,916

and U.S. Patent No. 4,513,006, which also disclose processes for the preparation of said compounds; and which are hereby incorporated by reference.

One process disclosed in the above referenced patents is a process for the preparation of the compounds of formula (I) comprising reacting an alcohol of the formula RCH_2OH with a chlorosulfamate of the formula $ClSO_2NH_2$ or $ClSO_2NHR^1$ in the presence of a base such as potassium t-butoxide or sodium hydride at a temperature of about $-20^\circ C$ to $25^\circ C$ and in a solvent such as toluene, tetrahydrofuran or dimethylformamide, where R is a moiety of the formula (II)



This process has two major disadvantages. One disadvantage is that the process calls for a combination of NaH and DMF which has an uncontrollable exotherm and is therefore potentially explosive. See J. Buckley et al., Chemical & Engineering News, July 12, 1982, page 5; and G. DeWail, Chemical & Engineering News, Sept. 13, 1982. Another disadvantage is that the process also uses highly toxic and corrosive chlorosulfonyl isocyanate (CSI) to prepare the commercially unavailable sulfamyl chloride ($ClSO_2NH_2$). The CSI is not only difficult to work with, because of its toxicity and corrosiveness, but also is available from only limited suppliers.

Another process for the preparation of compound of

formula (I) disclosed in the above mentioned U.S. Pat. No. 4,513,006 comprises reacting an alcohol of the formula RCH_2OH with sulfuryl chloride of the formula SO_2Cl_2 in the presence of a base such as triethylamine or pyridine at a temperature of about -40°C to 25°C in a diethyl ether or methylene chloride solvent to produce a chlorosulfate of the formula $\text{RCH}_2\text{OSO}_2\text{Cl}_2$. The chlorosulfate of the formula $\text{RCH}_2\text{OSO}_2\text{Cl}_2$ may then be reacted with an amine of the formula R^1NH_2 at a temperature of about -40°C to 25°C in a methylene chloride or acetonitrile solvent to produce the compound of formula (I). This process utilizing diethyl ether, methylene chloride and acetonitrile solvents produces relatively low yields of the desired end product of formula (I).

A third process disclosed in the two patents mentioned above comprises reacting the chlorosulfate of formula $\text{RCH}_2\text{OSO}_2\text{Cl}_2$ formed as previously described with a metal azide such as sodium azide in a solvent such as methylene chloride or acetonitrile to yield an azidosulfate of the formula $\text{RCH}_2\text{OSO}_3\text{N}_3$. The azidosulfate is then reduced to the compound of formula I wherein R^1 is hydrogen, by catalytic hydrogenation.

The disadvantage with this process is that explosions may occur when handling the azide compounds. Also, the process contains an additional chemical transformation involving the reduction of the azide to the NH_2 moiety.

Maryanoff et al. in US 5,387,700 disclose a process for the preparation of compounds of formula (I) which comprises reacting an alcohol of the formula RCH_2OH with sulfuryl chloride in the presence of a base, in a solvent


selected from the group consisting of toluene, t-butyl methyl ether and tetrahydrofuran, to form a chlorosulfate intermediate of the formula $RCH_2OSO_2Cl_2$. The chlorosulfate of formula $RCH_2OSO_2Cl_2$ is reacted with an amine of the
5 formula R_1NH_2 , in a solvent selected from the group consisting of tetrahydrofuran, t-butyl methyl ether and lower alkanol (e.g. methanol or ethanol) to form the compound of formula (I).

One disadvantage of this process is that the compound
10 of formula (I) is prepared in a batch process wherein the first reaction is carried out, the solvent is removed, the product is isolated, the isolated product is re-dissolved in a second solvent and then reacted to form the final product. This results in a process which requires
15 isolation of a semi-stable, thermally labile $ROSO_2Cl$ intermediate. Additionally, this process requires handling of multiple solvents and multiple solvent recovery processes for recycling or disposal resulting in a process which is both cost and labor intensive.

It is an object of the present invention to provide a
20 one step process for the preparation of the compounds of formula (I), a process which uses readily available materials, can be carried out without isolation of
25 intermediates and/or produces relative high yields, thereby allowing for commercial production of the compounds of formula (I).

Summary of the Invention

30 The present invention is directed to a one-step process for the preparation of the compounds of formula (I)




wherein

X is selected from CH₂ or O;

R¹ is selected from the group consisting of hydrogen
5 and C₁₋₄alkyl;


R³, R⁴, R⁵ and R⁶ are each independently selected from
hydrogen or lower alkyl and, when X is CH₂, R⁵ and R⁶ may
be alkene groups joined to form a benzene ring and, when X
is O, R³ and R⁴ and/or R⁵ and R⁶ together may be a
10 methylenedioxy group of the formula:



wherein

R⁷ and R⁸ are same or different and are hydrogen,
lower alkyl or are alkyl and are joined to form a
15 cyclopentyl or cyclohexyl ring;


comprising




reacting a compound of formula (II) with sulfonyl
diamide (also known as sulfamide), at an elevated
20 temperature, in the presence of from 0 to about 10% water,
to yield the corresponding compound of formula (I).

The present invention is further directed to a

process for the preparation of a compound of formula (Ia),



also known as topiramate, comprising



5 reacting a compound of formula (IIa) (also known as diacetone fructose or DAF) with sulfuryl diamide (also known as sulfamide), at an elevated temperature, in the presence of from 0 to about 10% water, to yield the corresponding compound of formula (Ia).

10 The present invention is further directed to a compound prepared according to any of the processes described herein.

15 Illustrative of the invention is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound prepared according to any of the processes described above.


20 An illustration of the invention is a pharmaceutical composition made by mixing a pharmaceutically acceptable carrier and a compound prepared according to any of the processes described above.

Illustrating the invention is a process for making a pharmaceutical composition comprising mixing a pharmaceutically acceptable carrier and a compound prepared according to the processes described above.


Another example of the invention is the use of a compound prepared according to any of the processes described herein in the preparation of a medicament for treating epilepsy.

Detailed Description of the Invention

In an embodiment of the present invention, the compound of formula (I) is topiramate, a compound of the formula (Ia)



As used herein, the terms "diacetone fructose" and "DAF" shall mean diacetone- β -D-fructose, a compound of formula (IIa)



As used herein, the term "alkyl", whether used alone

or as part of a substituent group, shall include straight and branched carbon chains. For example, alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, and the like. Unless
5 otherwise noted, the term "**lower**" when used with alkyl shall mean carbon chain compositions of one to four carbon atoms.

As used herein, the notation "*" shall denote the
10 presence of a stereogenic center.

As used herein, the term "**elevated temperature**" shall mean a temperature greater than about 90°C, preferably a temperature greater than or equal to the reflux
15 temperature of the solvent of the reaction mixture, more preferably a temperature in the range of about 90°C to about 170°C, more preferably still at a temperature in the range of about 110°C to about 160°C, more preferably still at a temperature in the range of about 120°C to about
20 140°C.

As used herein, unless otherwise noted, the term "**in the presence of from 0 to about 10% water**" shall mean that the total molar amount of water relative to the molar
25 amount of DAF is in the range of from 0 to about 10%.

As used herein, the term "**aprotic organic solvent**" shall mean any organic solvent that does not yield a proton under the reaction conditions. Suitable examples
30 include, but are not limited to, xylene (for example o-xylene, p-xylene, m-xylene or a mixture thereof), ethyl

benzene, mesitylene, tetrahydronaphthalene, pyridine, 1-methyl-2-pyrrolidinone, toluene, 4-methyl-2-pentanone, benzonitrile, dimethylformamide, sulfolane, diphenyl ether, and the like.

5

The term "**subject**" as used herein, refers to an animal, preferably a mammal, more preferably a human, who is or has been the object of treatment, observation or experiment.

10

The term "**therapeutically effective amount**" as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical

15


doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term "**composition**" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

20


25

One skilled in the art will recognize that the compounds of formula (I) have several stereogenic centers as denoted by the asterisks below.



One skilled in the art will further recognize that the processes of the present invention may be used to prepare racemic mixtures of a compound of the formula (I) or any of the stereoisomers of a compound of formula (I),
 5 by selection and substitution of appropriate racemic mixtures or stereoisomers of the reagents.

The present invention is directed to a one step process for the preparation of a compound of formula (I)
 10 as outlined in Scheme 1.



Scheme 1

Accordingly, a suitably substituted compound of
 15 formula (II), a known compound or compound prepared by known methods, is reacted with sulfuryl diamide (also known as sulfamide),

wherein the sulfuryl diamide is preferably present in an amount greater than or equal to about 0.9 equivalents; more preferably in an amount in the range of about 1.5 to
 20 about 3 equivalents; more preferably still in an amount equal to about 2 equivalents;

preferably in the presence of a non-aqueous organic base such as an organic tertiary amine such as pyridine, 4-picoline, isoquinoline, dimethylbutylamine, 4-
 25 dimethylaminopyridine, 4-t-butylpyridine, imidazole, tributylamine, dimethylbenzylamine, and the like; or a non-aqueous inorganic base such as K_2CO_3 , KHCO_3 , Na_2CO_3 , NaHCO_3 , and the like; more preferably, in the presence of a

non-aqueous organic tertiary amine; more preferably still, in the presence of pyridine;

wherein the non-aqueous organic or inorganic base is preferably present in an amount greater than about 1
5 equivalent; more preferably, in an amount in the range of about 3 to about 6 equivalents; more preferably still, in an amount equal to about 4 equivalents;

in the presence of from 0% to about 10% water, preferably from 0% to about 5% water, more preferably from
10 0% to about 3% water;

preferably in an aprotic organic solvent such as xylene (for example *o*-xylene, *p*-xylene, *m*-xylene or a mixture thereof), ethyl benzene, mesitylene, tetrahydronaphthalene, pyridine, 1-methyl-2-pyrrolidinone,
15 toluene, 4-methyl-2-pentanone, benzonitrile, dimethylformamide, sulfolane, diphenyl ether, and the like; or a mixture of aprotic organic solvents such as diphenyl ether:biphenyl, xylene:toluene, technical grade mixed xylenes, and the like; more preferably in *o*-xylene,
20 *p*-xylene, *m*-xylene, a mixture thereof or a technical grade mixed xylenes;

alternatively, the solvent may be selected from a non-aqueous organic base such as an organic tertiary amine, for example, pyridine, 4-picoline, isoquinoline,
25 dimethylbutylamine, 4-dimethylaminopyridine, 4-*t*-butylpyridine, imidazole, tributylamine, dimethylbenzylamine, and the like; preferably, a non-aqueous organic tertiary amine; more preferably, pyridine;


alternatively still, when the compound of formula
30 (II) is reacted in the presence of a non-aqueous organic base, the non-aqueous organic base may act as the solvent; at an elevated temperature, preferably at a

temperature in the range of from about 90°C to about 170°C;
more preferably, at a temperature in the range of from
about 110°C to about 160°C; more preferably still at a
temperature in the range of from about 120°C to about
5 140°C;

to yield the corresponding compound of formula (I).

Preferably, the compound of formula (II) is reacted
with sulfuryl diamide, according to the process outlined
10 in Scheme 1 above, in the absence of any other alcohol
(i.e. an alcohol other than the compound of formula (II)).

The compound of formula (Ia) may be prepared
according to the process outlined in Scheme 2.



Scheme 2

Accordingly, a compound of formula (IIa), a known
compound also known as diacetone fructose (DAF), is
reacted with sulfuryl diamide (also known as sulfamide),

20 wherein the sulfuryl diamide is preferably present in
an amount greater than or equal to about 0.9 equivalents;
more preferably in an amount in the range of about 1.5 to
about 3 equivalents; more preferably still in an amount
equal to about 2 equivalents;

25 preferably in the presence of a non-aqueous organic
base such as an organic tertiary amine such as pyridine,
4-picoline, isoquinoline, dimethylbutylamine, 4-

dimethylaminopyridine, 4-t-butylpyridine, imidazole, tributylamine, dimethylbenzylamine, and the like; or a non-aqueous inorganic base such as K_2CO_3 , $KHCO_3$, Na_2CO_3 , $NaHCO_3$, and the like; more preferably, in the presence of a non-aqueous organic tertiary amine; more preferably still, in the presence of pyridine;

wherein the non-aqueous organic or inorganic base is preferably present in an amount greater than about 1 equivalent; more preferably, in an amount in the range of about 3 to about 6 equivalents; more preferably still, in an amount equal to about 4 equivalents;

in the presence of from 0% to about 10% water, preferably from 0% to about 5% water, more preferably from 0% to about 3% water;

preferably in an aprotic organic solvent such as xylene (for example *o*-xylene, *p*-xylene, *m*-xylene or a mixture thereof), ethyl benzene, mesitylene, tetrahydronaphthalene, pyridine, 1-methyl-2-pyrrolidinone, toluene, 4-methyl-2-pentanone, benzonitrile, dimethylformamide, sulfolane, diphenyl ether, and the like; or a mixture of aprotic organic solvents such as diphenyl ether:biphenyl, xylene:toluene, technical grade mixed xylenes, and the like; more preferably in *o*-xylene, *p*-xylene, *m*-xylene, a mixture thereof or a technical grade mixed xylenes;

alternatively, the solvent may be selected from a non-aqueous organic base such as an organic tertiary amine, for example, pyridine, 4-picoline, isoquinoline, dimethylbutylamine, 4-dimethylaminopyridine, 4-t-butylpyridine, imidazole, tributylamine, dimethylbenzylamine, and the like; preferably, a non-aqueous organic tertiary amine; more preferably, pyridine;

alternatively still, when the compound of formula (II) is reacted in the presence of a non-aqueous organic base, the non-aqueous organic base may act as the solvent;

at an elevated temperature, preferably at a temperature in the range of from about 90°C to about 170°C; more preferably, at a temperature in the range of from about 110°C to about 160°C; more preferably still at a temperature in the range of from about 120°C to about 140°C;

to yield the corresponding compound of formula (Ia).

Preferably, the compound of formula (IIa) is reacted with sulfuryl diamide, according to the process outlined in Scheme 2 above, in the absence of any other alcohol (i.e. an alcohol other than the compound of formula (IIa)).

Where the compounds of the present invention have at least one chiral (or stereogenic) center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral (or stereogenic) centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention.

Where the processes for the preparation of the compounds according to the invention give rise to a mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative
5 chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation
10 of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-D-tartaric acid and/or (+)-di-p-toluoyl-L-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of
15 diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

20 During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those
25 described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991. The protecting groups may be removed
30 at a convenient subsequent stage using methods known from the art.

The following examples describe the invention in

greater detail and are intended to illustrate the invention, but not to limit it.

Example A1

5 **Experimental Runs #101-116:**

 A reactor was charged with DAF, sulfamide, base, and solvent, as described in Table 1. While agitating, the reaction mixture was heated and held at the desired temperature and for the selected amount of time (as listed
10 in Table 2).

 For experimental runs 101, 102, 103, the mixture was distilled at 41°C, under vacuum, for 30, 35 and 20 minutes, respectively.

 The reaction mixtures were cooled to room
15 temperature, then extracted twice with about 50 mL of a 1M aqueous NaOH solution (An 0.3 M solution of aqueous NaOH was used for experimental runs #101-108). The resulting basic product solution was neutralized using 6M aqueous HCl solution. After crystallization, the product solution
20 was filtered, and the wet cake was dried for between 16-20 hours (about 22 hours for experimental runs # 106 and 107) at about 50°C under vacuum to yield topiramate product. The dry product was analyzed by HPLC.

25 **Experimental Runs # 117-139:**

 A reactor was charged with DAF, sulfamide, base, and solvent, as described in Table 1. While agitating, the reaction mixture was heated and held for a period of about 60 minutes at the desired temperature (as listed in Table
30 2). After cooling to room temperature, the reaction mixture was extracted twice with about 75 mL of 1M aqueous NaOH solution. The resulting basic product solution was

neutralized using 6M aqueous HCl solution. After crystallization, the product solution was filtered, and the wet cake was dried for 16-20 hours (~24 hours for experimental runs 129, 134, 137 and ~48 hours for experimental runs # 126, 128, 130, 131, 136) at about 50°C with vacuum to yield topiramate product. Experimental runs # 123, 125 and 135 were expected to yield product, but were abandoned due to difficulties during extraction or crystallization. The dry product was analyzed by HPLC.

The above described procedures were completed according to the run conditions as listed in Table 1 and 2 below. Topiramate product was produced in all experimental runs except as noted above.

Table 1

Run #	DAF	Sulfamide	Base	Base	Solvent
101	10.05 g	7.43 g	pyridine	12.14 g	mixed xylenes
	38.62 mmol	77.32 mmol		153.48 mmol	100 mL
102	10.0 g	7.39 g	pyridine	12.16 g	mixed xylenes
	38.43 mmol	76.9 mmol		153.73 mmol	100 mL
103	10.0 g	7.33 g	pyridine	12.18 g	mixed xylenes
	38.43 mmol	76.27 mmol		153.98 mmol	100 mL
104	10.0 g	7.43 g	pyridine	12.34 g	mixed xylenes
	38.43 mmol	77.32 mmol		156.01 mol	100 mL
105	10.01 g	7.39 g	pyridine	12.17 g	mixed xylenes
	38.47 mmol	76.9 mmol		153.86 mmol	101 mL

106	10.01 g	7.39 g	pyridine	12.45 g	mixed xylenes
	38.47 mmol	76.9 mmol		157.4 mmol	101 mL
107	10.01 g	7.44 g	pyridine	12.15 g	mixed xylenes
	38.47 mmol	77.42 mmol		153.6 mmol	51 mL
108	10.02 g	7.38 g	4- picoline	12.54 g	mesitylene
	38.50 mmol	76.8 mmol		134.65 mmol	50 mL
109	14.98 g	11.08 g	pyridine	18.33 g	mixed xylenes
	57.56 mmol	115.3 mmol		231.73 mmol	150 mL
110	9.99 g	7.39 g	Pyridine	12.14 g	O-xylene
	38.39 mmol	76.9 mmol		153.48 mmol	99 mL
111	10.04 g	7.31 g	Pyridine	12.14 g	O-xylene
	38.58 mmol	76.07 mmol		153.48 mmol	100 mL
112	10.03 g	7.39 g	Pyridine	12.22 g	O-xylene
	38.54 mmol	76.9 mmol		154.49 mmol	100 mL
113	10.02 g	7.39 g	Pyridine	12.22 g	O-xylene
	38.50 mmol	76.9 mmol		154.49 mmol	100 mL
114	9.98 g	7.38 g	Pyridine	12.25 g	O-xylene
	38.35 mmol	76.8 mmol		154.87 mmol	100 mL
115	10.02 g	7.36 g	pyridine	12.4 g	O-xylene
	38.5 mmol	76.59 mmol		156.76 mmol	99 mL
116	10.0 g	7.36 g	Pyridine	12.16 g	O-xylene
	38.43 mmol	76.59 mmol		153.73 mmol	100 mL
117	29.92 g	22.31 g	Pyridine	36.52 g	O-xylene
	114.98 mmol	232.15 mmol		461.49 mmol	302 mL

118	30.06 g	22.3 g	Pyridine	36.68 g	O-xylene
	115.51 mmol	232.05 mmol		463.72 mmol	150 mL
119	30.07 g	22.34 g	4- picoline	42.48 g	1:1 O- xylene: 4- picoline
	115.55 mmol	232.47 mmol		456.14 mmol	150 mL
120	30.03 g	22.22 g	Pyridine	36.58 g	O-xylene
	115.4 mmol	231.22 mmol		462.45 mmol	75 mL
121	30.04 g	22.23 g	Pyridine	36.55 g	O-xylene
	115.44 mmol	231.32 mmol		462.07 mmol	40.0 mL
122	29.98 g	22.18 g	pyridine	36.37 g	none
	115.09 mmol	230.8 mmol		459.8 mmol	
123	30.02 g	22.34 g	4- picoline	42.91 g	none
	115.36 mmol	232.47 mmol		460.75 mmol	
124	30.09 g	22.24 g	iso- quinoline	58.11 g	none
	115.63 mmol	231.43 mmol		449.91 mmol	
125	30.01 g	22.18 g	4- picoline	42.92 g	tetrahydro- naphthalene
	115.32 mmol	230.8 mmol		460.86 mmol	40 mL

126	30.01 g	22.17 g	4- picoline	42.8 g	tetrahydro- naphthalene
	115.32 mmol	230.7 mmol		459.57 mmol	41 mL
127	29.99 g	22.34 g	dimethy- butyl- amine	59.95 g	tetrahydro- naphthalene
	115.24 mmol	232.47 mmol		757.9 mmol	40 mL
128	30.02 g	22.22 g	pyridine	36.56 g	tetrahydro- naphthalene
	115.36 mmol	231.22 mmol		462.2 mmol	40 mL
129	29.95 g	22.32 g	pyridine	36.45 g	diphenyl ether
	115.09 mmol	232.26 mmol		460.81 mmol	41 mL
130	30.02 g	22.34 g	4- picoline	43.03 g	diphenyl ether
	115.36 mmol	232.47 mmol		462.04 mmol	40.5 mL
131	29.99 g	22.24 g	4- picoline	42.94 g	O-xylene
	115.24 mmol	231.43 mmol		461.08 mmol	40 mL
132	30.03 g	22.5 g	pyridine	36.12 g	O-xylene
	115.4 mmol	234.31 mmol		456.64 mmol	40 mL
133	10.04 g	7.38 g	iso- quinoline	20.05 g	O-xylene
	38.58 mmol	76.8 mmol		155.23 mmol	50 mL

134	30.05 g	22.21 g	4-t-butyl-pyridine	62.3 g	O-xylene
	115.47 mmol	231.11 mmol		460.76 mmol	40 mL
135	30.08 g	22.45 g	iso-quinoline	59.65 g	diphenyl ether
	115.59 mmol	233.61 mol		461.83 mmol	40.5 mL
136	60.08 g	44.28 g	4-picoline	85.91 g	O-xylene
	230.87 mmol	460.77 mmol		922.47 mmol	75 mL
137	59.99 g	44.44 g	Pyridine	73.04 g	O-xylene
	230.53 mmol	462.43 mmol		923.39 mmol	75 mL
138	30.04 g	22.5 g	4-t-butyl-pyridine	46.66 g	O-xylene
	115.44 mmol	234.13 mmol		345.09 mmol	75 mL
139	30.02 g	22.18 g	4-t-butyl-pyridine	46.68 g	O-xylene
	115.36 mmol	230.8 mmol		345.25 mmol	75.5 mL

Table 2

Run #	Time (min)	Temp. (°C)
101	~65	128-140
102	60	135-136

103	65	132-136
104	60	138-139
105	60	134-137
106	65	130-136
107	60	124-133
108	Not available	130-140
109	80	99-114
110	55	133-135
111	45	130-133
112	50	130-134
113	60	135-137
114	60	134-136
115	62	133-137
116	60	134-135
117	62	135-139
118	60	134-136
119	62	141-146
120	62	130-135
121	60	131-133
122	69	132-134
123	60	150-156
124	62	142-144
125	62	153-157
126	55	130-140
127	60	109-112
128	61	137-141
129	58	139-145
130	67	136-140
131	62	134-144
132	60	130-138
133	60	140-135

134	64	133-132
135	57	130-134
136	64	130-138
137	60	130-132
138	34	130-149
139	41	25-120

Example A2

Experimental Runs #140-147; Pyridine Base:

Eight experiments were performed to evaluate combined
5 effects of solvent volume, reaction time, and reaction
temperature on one-step synthesis of topiramate, starting
with diacetone fructose (DAF).

DAF was combined with sulfamide, pyridine and the
10 chosen volume of o-xylene solvent (in amounts as listed in
Table 3) in a 1 liter round-bottomed flask. The charged
flask was then submerged in a pre-heated oil bath at the
chosen external temperature, and the reaction mixture
agitated and heated to the chosen reaction temperature,
15 where it was held for the chosen time before the flask was
removed from the oil bath and cooled to room temperature.

The reaction mixture was then extracted twice with 1M
NaOH (75mL portions). The aqueous extracts were combined
and neutralized with 6M HCl to induce product
20 crystallization. The product precipitated as an oil in
all cases. The oil was agitated overnight to become
filterable crystals. The resulting solids were dried
under vacuum for 16-18 hrs at 50°C. The dried product was
analyzed by HPLC.

25 Following the above procedure the eight experiments
were performed with experimental conditions as listed in

Table 3 and 4 below. All experimental runs yielded topiramate product.

Table 3

Pyridine Base				
Run #	DAF	Sulfamide	Base	o-Xylene
140	30.01 g	22.24 g	36.56 g	75 mL
	115.32 mmol	231.43 mmol	462.2 mmol	
141	30.03 g	22.37 g	36.78 g	75.5 mL
	115.4 mmol	232.78 mmol	464.98 mmol	
142	30.01 g	22.58 g	36.5 g	75.5 mL
	115.32 mmol	234.96 mmol	461.44 mmol	
143	30.12 g	22.44 g	36.68 g	75 mL
	115.74 mmol	233.51 mmol	463.72 mmol	
144	30 g	22.34 g	36.88 g	40 mL
	115.28 mmol	232.47 mmol	466.25 mmol	
145	30.05 g	22.31 g	36.51 g	40 mL
	115.47 mmol	232.15 mmol	461.57 mmol	
146	30.03 g	22.24 g	36.6 g	40 mL
	115.4 mmol	231.43 mmol	462.71 mmol	
147	30.07 g	22.34 g	36.61 g	40 mL
	115.55 mmol	232.47 mmol	462.83 mmol	

5

Table 4

Run #	Time (min)	Temp (°C)
140	60	130.0-133.4
141	30	130.0-132.7
142	60	130.0-134.3
143	30	130.0-134.7
144	30	130.0-130.3
145	60	130.0-134.9

146	60	130.0-132.8
147	30	130.0-133.9

Example A3

Experimental Runs #148-156: t-Butylpyridine Base:

Nine experiments were performed to evaluate the combined effects of base equivalents, reaction temperature and reaction time on one-step synthesis of topiramate using 4-tert-butylpyridine and starting with diacetone fructose (DAF).

DAF was combined with sulfamide, 4-tert-butylpyridine (in amounts as listed in Table 4) and 40 mL o-xylene in a 500 mL round-bottomed flask. The reaction mixture was agitated and heated to the chosen reaction temperature in an oil bath over about 19-23 min. The reaction mixture was maintained at the chosen reaction temperature for the chosen time, and then cooled rapidly to 25°C. The reaction mixture was extracted twice with 1M NaOH (75mL portions). The aqueous extracts were combined and neutralized using 6M HCl to induce product crystallization. The product precipitated as an oil in all cases, which was agitated overnight to become filterable crystals. The crystals were dried under vacuum for 16-18 hrs at 50°C. The dried product was analyzed by HPLC.

Following the above procedure the nine experiments were performed with experimental conditions as listed in Table 5 and 6 below. All experimental runs yielded topiramate product.

Table 5

4-t-butylpyridine Base

Run #	DAF	Sulfamide	Base	o-Xylene
148	30.16 g	22.42 g	47.07 g	40 mL
	115.9 mmol	231.4 mmol	348.1 mmol	
149	30.20 g	22.21 g	15.67 g	40 mL
	116.05 mmol	231.1 mmol	115.9 mmol	
150	30.12 g	22.16 g	47.03 g	40 mL
	115.7 mmol	230.6 mmol	347.8 mmol	
151	30.23 g	22.12 g	31.19 g	40 mL
	116.2 mmol	230.2 mmol	231.7 mmol	
152	30.10 g	22.75 g	46.76 g	40 mL
	115.7 mmol	236.8 mmol	345.8 mmol	
153	30.03 g	22.42 g	15.76 g	40 mL
	115.4 mmol	233.3 mmol	116.6 mmol	
154	30.23 g	22.24 g	46.83 g	40 mL
	116.2 mmol	231.5 mmol	346.4 mol	
155	30.15 g	22.49 g	46.85 g	40 mL
	115.9 mmol	234.1 mmol	346. mmol	
156	30.08 g	22.24 g	15.60 g	40 mL
	115.6 mmol	231.5 mmol	115.4 mmol	

Table 6

Run #	Time (min)	Temp (°C)
148	44	130.0-150.5
149	60	110.0-126.6
150	60	110.0-129.0
151	28	130.1-146.6
152	29	130.0-151.6
153	15	130.0-141.5
154	30	130.1-143.4
155	15	110.1-117.4
156	30	110.1-113.7

Example A4

Multiple experiments were run to evaluate the effect of solvent and base on the one step process for the preparation of topiramate. Experimental conditions were as listed in Tables 7 and 8, with experimental procedures for the individual runs as described below.

Run #157, 158, 160, 164-166:

DAF and sulfamide (in amounts as in Table 8) were transferred to a test tube. The chosen solvent and the chosen base (in amounts as listed in Table 7) were then added to the DAF/sulfamide mixture. The reaction mixture was then agitated and heated, with the reaction mixture maintained at the selected temperature (see Table 8) for the selected time (see Table 8), after which time the upper layer was sampled and analyzed by HPLC.

Run #159,188-198:

DAF, sulfamide, o-xylene and the specified amount of the chosen base (see Tables 7 and 8) were transferred into a 500mL round-bottomed flask. The reaction mixture was then agitated and heated, with the mixture maintained at the chosen temperature for the chosen time (see Table 8) with an oil bath. The reaction mixture was removed from the hot oil, cooled to about room temperature and then extracted twice with 1M NaOH (75 mL portions). The aqueous extracts were combined and neutralized using 6M HCl to induce product crystallization. The resulting solids were dried under vacuum for 16-18 hrs at 50°C. The dried product was analyzed by HPLC.

Run #161-163

DAF, sulfamide, 4-picoline, and the chosen solvent (in amounts as in Table 7 and 8) were transferred into a 250mL round-bottomed flask. The flask was lowered into a heated oil bath and the mixture agitated at the selected holding temperature for the selected reaction time (see Table 8). The reaction mixture was then cooled to room temperature and extracted twice with 1M NaOH (75mL portions). The aqueous extracts were neutralized with 6M HCl and then maintained at room temperature overnight to allow product to crystallize. The resulting crystalline solids were filtered and dried for 16-18 hrs in a vacuum oven at 50°C. The product was then analyzed by HPLC.

Note: Although product was expected in run #162, the product did not crystallize and was therefore not analyzed; the product from run #163 was a sticky solid.

Run #167-172:

The selected solvent in the selected amount (see Table 7) was transferred to a 125 mL test tube along pyridine (in amounts as in Table 7) and the resulting solution heated to the chosen temperature. A pre-weighed mixture of DAF and sulfamide (in amounts as in Table 8) was then added along with transfer washes of solvent (about 2 mL). The reaction mixture was agitated at the chosen temperature for the chosen time (see Table 8) at which time the upper layer was sampled (some mixtures formed a small, dark second layer) and analyzed by HPLC.

Run #173

DAF, sulfamide, pyridine and mixed xylenes (in amounts as listed in Tables 7 and 8) were transferred to a

1000 mL round-bottomed flask in a pre-heated (140°C) oil bath. The reaction mixture was agitated at the selected reaction temperature for 90 mins. The flask was removed from the oil bath and the residual heat used to distill about 22g of liquid from the reaction mixture. The reaction mixture was extracted with three portions (90mL, 30mL, and 10mL) of 0.35M NaOH. 1N HCl (50mL) was added to the aqueous extracts, resulting in the formation of an oil. The mixture was agitated to transform the oil into filterable crystals. The crystals were collected by filtration, washed with water, and dried in a vacuum oven for 16 hrs at 50°C. The product was then analyzed by HPLC analysis.

Run #174-180:

DAF, sulfamide, pyridine and the chosen solvent (in amounts as listed in Tables 7 and 8) were transferred to a 500 ml round-bottomed flask. The reaction mixture was then agitated and heated in an oil bath to the target temperature. The reaction mixture was maintained at about the target temperature (listed in Table 8) for the selected time (see Table 8), removed from the hot oil and cooled to about room temperature. To the reaction mixture was then added water (100mL) water and/or other solvents (MeOH, 0.5M NaOH, or the reaction solvent) as needed, to achieve a heterogeneous mixture consisting of two clear liquid layers. GC analysis was then performed on both layers.

Run# 181-183

DAF, sulfamide, 4-tert-butylpyridine, and the selected solvent (in amounts as listed in Tables 7 and 8)

were transferred to a 500 mL round-bottomed flask. The reaction mixture was then placed in oil bath and agitated while heating at the selected reaction temperature for the selected time (see Table 8). The reaction mixture was then allowed to cool to room temperature. The reaction mixture was then extracted with the appropriate solvents to yield two clear liquid layers (For run 181 the extraction solvents were water (125mL) and mixed xylenes (25 mL); for run 182 the extraction solvents were water (150mL) and methanol (25mL); for run 183 the extraction solvent was 0.3M NaOH (100mL)). GC analysis was performed on liquid samples from both layers.

Run #184

DAF, sulfamide, dimethylbutylamine and 1,2,3,4-tetrahydronaphthalene (in amounts as listed in Tables 7 and 8) were transferred into a 250mL round-bottomed flask. The flask was placed in a heated oil bath (136°C) and the mixture agitated while heating at the selected temperature for the selected time (see Table 8). The reaction mixture was cooled to about room temperature and then extracted twice with 1M NaOH (75mL portions). The aqueous extracts were neutralized using 6M HCl and agitated for 2hrs to allow crystallization. The crystals were collected by filtration and then dried for about 17 hrs 45 min in a vacuum oven at 52°C. The product was then tested by HPLC analysis.

Run #185-187

DAF, sulfamide, isoquinoline and the chosen solvent (in amounts as listed in Tables 7 and 8) were transferred to a 250 mL round-bottomed flask. The reaction mixture

was then placed in an oil bath and agitated while heating at the selected reaction temperature for the selected time (see Table 8). The reaction mixture was cooled to about room temperature and then extracted twice with 1M NaOH (75mL portions). The aqueous extracts were neutralized with 6M HCl and allowed to stand at room temperature for several days to facilitate crystallization. The crystals were collected by filtration and then dried about 16-18hrs in a vacuum oven at 50°C. The dried product was analyzed by HPLC analysis.

Note: All experimental runs yielded topiramate product except as noted herein. An emulsion was observed to form during the extraction of Experimental Run #187 and although topiramate product was expected, no product was collected. The product collected from Experimental Runs #187, 191, 194 and 197 was expected to be topiramate, but not analyzed.

Tables 7 and 8 below list the run conditions for the experimental procedures described above. Except as noted above, all experimental runs yielded topiramate product.

Table 7

Run#	Base	Base (g)	Solvent	Solvent (mL)
157	Imidazole	1.01	p-Xylene	15.1
158	4-(Dimethylamino)-pyridine	1.77	p-Xylene	15.2
159	4-(Dimethylamino)-pyridine	42.08	o-Xylene	40
160	4-Picoline	1.38	p-Xylene	15
161	4-Picoline	42.48	o-Xylene	150

162	4-Picoline	42.92	Tetrahydro-napthalene	40
163	4-Picoline	43.03	Diphenyl Ether	40.5
164	2,6-Lutidine	1.55	p-Xylene	15
165	Pyridine	1.15	p-Xylene	15
166	Pyridine	14.7	pyridine	15
167	Pyridine	1.17	1-Methyl-2-pyrrolidinone	15
168	Pyridine	1.14	Toluene	15
169	Pyridine	1.15	4-Methyl-2-pentanone	15
170	Pyridine	1.15	Benzonitrile	15
171	Pyridine	1.15	p-Xylene	15.1
172	Pyridine	1.17	Dimethyl-formamide	15
173	Pyridine	17.29	mixed xylenes	300
174	Pyridine	18.23	Mesitylene	150
175	Pyridine	18.22	o-Xylene	150
176	Pyridine	18.22	Tetramethylene sulfone	150
177	Pyridine	18.22	Biphenyl	150g
178	Pyridine	18.20	Tetrahydro-napthalene	150
179	Pyridine	18.23	Diphenyl Ether	151
180	Pyridine	12.17	3:1 Diphenyl Ether: Biphenyl	100.04g
181	4-tert-Butylpyridine	31.17	mixed xylenes	150
182	4-tert-Butylpyridine	----	4-tert-Butylpyridine	75
183	4-tert-Butylpyridine	20.78	mesitylene	100

184	Dimethylbutylamine	59.95	Tetrahydro-napthalene	40
185	isoquinoline	58.11	isoquinoline	-----
186	isoquinoline	20.05	o-Xylene	50
187	isoquinoline	59.65	Diphenyl Ether	40.5
188	Tributylamine	85.49	o-Xylene	40
189	Dimethyl-benzylamine	62.33	o-Xylene	40
190	Dimethyldodecyl-amine	98.42	o-Xylene	40
191	Hexadecylamine	83.62	o-Xylene	40
192	Dimethyltetradecyl-amine	111.22	o-Xylene	40
193	Dimethylhexadecyl-amine	124.30	o-Xylene	40
194	Dimethyloctadecyl-amine	137.13	o-Xylene	40
195	morpholine	30.30	o-Xylene	40
196	pyrrolidine	25.74	o-Xylene	40
197	urea	20.29	o-Xylene	40
198	(none)	---	o-Xylene	40

Table 8

Run #	DAF (g)	Sulfamide (g)	Time (min)	Target Temp (C)
157	0.76	0.84	30	115
158	0.74	0.84	30	115
159	30.21	22.21	30	130
160	0.76	0.84	30	115
161	30.07	22.34	62	140
162	30.01	22.18	62	150

163	30.02	22.34	67	140
164	0.75	0.84	30	115
165	0.76	0.84	30	115
166	0.74	0.84	30	115
167	0.73	0.83	31	115
168	0.75	0.85	30	115
169	0.75	0.84	30	115
170	0.76	0.85	30	115
171	0.77	0.85	30	115
172	0.76	0.83	30	115
173	15.02	19.46	90	130
174	15.02	11.08	60	130
175	15.05	11.07	60	130
176	15.01	11.08	60	130
177	15.04	11.08	60	130
178	15.02	11.08	60	130
179	15.03	11.07	60	130
180	10.03	7.38	60	140
181	15.03	11.08	60	130
182	7.49	5.53	60	130
183	10	7.38	60	150
184	29.99	22.34	60	110
185	30.09	22.24	62	140
186	10.04	7.38	60	140
187	30.08	22.45	57	130
188	30.05	22.31	30	130
189	30.03	22.22	30	140
190	30.17	22.26	30	130
191	30.09	22.04	30	130
192	30.1	22.21	30	130
193	30.11	22.28	30	130

194	30.18	22.33	30	130
195	30.06	22.26	30	130
196	30.01	22.26	30	130
197	30.13	22.32	30	130
198	30.02	22.16	30	130

Example A5

Topiramate was prepared according to the process of the present invention. Listed below are experimental procedures for the experiments. Tables 9 and 10 which follow list detailed experimental conditions.

Experiment 2:

DAF, sulfamide, pyridine, and o-xylene in amounts as listed in Table 9 below were charged to a 1-liter reactor.

The reactor was heated to a jacket remperature of about 155°C and maintained at that temperature for 30 minutes. The reactor was cooled to 60°C and distillation was started under vacuum, with distillation continued until there was no remaining liquid to distill. 1M NaOH was added to separate the reaction mixture into layers. The bottom, aqueous layer was drained and transferred to a separatory funnel. 1M NaOH was added to separate the layers further. Again, the lower, aqueous layer was drained and transferred to a clean reactor. 6M HCl was added to neutralize solution to pH 7. The product crystallized, was filtered, and then dried to yield topiramate.

Experiments 3-8, 19-21:

DAF, sulfamide, pyridine, and o-xylene in amounts as listed in Table 9 below were charged to a 1-liter reactor.

The reactor was heated and held at the selected temperature for the selected time (see Table 10), then cooled rapidly to 65°C. Once cooled, vacuum was applied, resulting in the distillation of some amount of organic liquids. The temperature during distillation was maintained between 55°C and 75°C. After distillation, some amount (0 to 200 milliliters) of NaOH (from strength of 1M to 3.5M) was added to split the reaction mixture into two separable phases, organic on top and aqueous on bottom. The bottom, aqueous layer was drained and transferred to a clean reactor. Acid (6M HCl) was added to neutralize the reaction mixture to pH 7. The product crystallized upon neutralization, was filtered and then dried to yield topiramate.

Experiment 9:

DAF, sulfamide, pyridine, and o-xylene in the amounts as listed in Table 9 below were charged to a 1-liter reactor. The reactor was heated to the selected temperature. The reactor was held at the maximum temperature for the selected amount of time (see Tables 9 and 10) and then cooled rapidly to 65°C. Once cooled, vacuum was applied, resulting in the distillation of some of the organic liquids. The temperature during distillation was maintained between 55°C and 75°C. After distillation, 3.5M NaOH was added to split the reaction mixture into two separable phases, organic on top and aqueous on bottom. The bottom, aqueous layer was drained and transferred to a clean reactor. The reactor was then spiked with pyridine (40 mL). Acid (6M HCl) was added to neutralize the reaction mixture to about pH 7. The product crystallized upon neutralization and was then

filtered and dried to yield topiramate.

Experiments 10, 15-17:

DAF, sulfamide, pyridine, and o-xylene in amounts as
5 listed in Table 9 below were charged to a 1-liter reactor.

The reactor was heated to the selected temperature. The
reactor was held at the selected temperature for the
selected time (see Tables 9 and 10) and then cooled
rapidly to 65°C. Once cooled, 3.5M NaOH was added to
10 split the reaction mixture into two separable phases,
organic on top and aqueous on bottom. The bottom, aqueous
layer was drained and transferred to a clean reactor.

Vacuum and heat were applied to distill off aqueous
liquid. Acid (6M HCl) was added to neutralize the
15 reaction mixture to about pH 7. If product did not
crystallize after neutralization, additional water was
added: for Experiment #15 350mL water was added; for
experiments #16 390 mL water was added; for Experiment #17
120 mL water was added. When the product had
20 crystallized, it was filtered and dried to yield
topiramate.

Experiment 11:

DAF, sulfamide, pyridine, and o-xylene in amounts as
25 listed in Table 9 below were charged to a 1-liter reactor.

The reactor was heated to the selected temperature, held
at this temperature for the selected time (see Tables 9
and 10) and then cooled rapidly to 65°C. Once cooled,
3.5M NaOH was added to split the reaction mixture into two
30 separable phases, organic on top and aqueous on bottom.
The bottom, aqueous layer was drained and transferred to a
clean reactor. Vacuum and heat were applied, resulting in

the distillation of some of the aqueous liquid. The reactor was heated to 60°C, and acid (6M HCl) was added to neutralize the reaction mixture to pH 7. The reactor was cooled to 1.5°C and then water was added until crystals
5 formed. The product crystallized and was then filtered and dried to yield topiramate.

Experiments 12-14:

DAF and sulfamide, in amounts as listed in Table 9
10 below were charged to a 1-liter reactor. The reactor was heated to the selected temperature. The reactor was held at the selected temperature for the selected time (see Tables 9 and 10) and then cooled rapidly to about room temperature. Once cooled, 3.5M NaOH was added to split
15 the reaction mixture into two separable phases, organic on top and aqueous on bottom. The bottom, aqueous layer was drained and transferred to a clean reactor. Acid (6M HCl) was added to neutralize the reaction mixture to about pH 7. The product crystallized upon neutralization and was
20 then filtered and dried to yield topiramate.

Experiment 18:

DAF, sulfamide, pyridine, and o-xylene in amounts as listed in Table 9 below were charged to a 1-liter reactor.
25 The reactor was heated to the selected temperature. The reactor was held at the selected temperature for the selected time (see Tables 9 and 10) and then cooled rapidly to 65°C. Once cooled, 3.5M NaOH was added to split the reaction mixture into two separable phases,
30 organic on top and aqueous on bottom. The bottom, aqueous layer was drained and transferred to a clean reactor. Vacuum and heat were applied to distill off about half of

the aqueous liquid. The remaining aqueous solution was drained and transferred to a beaker. 6M HCl (60 mL) was added to the cleaned reactor. Reaction mixture and additional 6M HCl were dosed into the reactor to maintain about pH 7. Water was added to induce crystallization. The product crystallized and was then filtered and dried to yield topiramate.

Experiment 19:

DAF, sulfamide and pyridine, in amounts as listed in Table 9 below were charged to a 1-liter reactor. The reactor was heated to the selected temperature. The reactor was held at the selected temperature for the selected time (see Tables 9 and 10) and then cooled rapidly. Once cooled to 22°C, 3.5M NaOH was added to split the reaction mixture into separable phases. Three fifths of the aqueous layer was drained and transferred to a clean reactor. Acid (6M HCl) was then added to neutralize the reaction mixture to about pH 7. Water 100 mL was added to the reaction mixture. The product crystallized, and was then filtered and dried to yield topiramate.

Table 9

Run#	DAF (gms) (mmols)	Sulfamide (gms) (mmols)	Pyridine (gms) (mmols)	O-xylene (mL)	Jacket Temp (°C)
2	29.97	22.26	36.50	40.0	155
	115.17	231.63	461.44		
3	100.01	73.86	121.40	134.0	145
	384.31	768.57	1534.77		
4	99.98	73.85	121.78	134.0	145

	384.20	768.47	1539.57		
5	100.04	78.21	121.82	133.0	145
	384.43	813.84	1540.08		
6	99.98	73.80	121.49	132.5	145
	384.20	767.95	1535.90		
7	99.99	73.93	121.66	134.0	145
	384.24	769.30	1538.05		
8	99.94	72.52	121.43	132.0	145
	384.04	754.63	1535.15		
9	100.01	73.50	12.11	134.0	145
	384.31	764.83	153.10		
10	100.01	73.86	121.84	132.0	145
	384.31	768.57	1540.33		
11	100.07	73.89	121.23	132.0	145
	384.54	768.89	1532.62		
12	100.01	73.87	0.00	0.0	145
	384.31	768.68	0.00		
13	100.02	73.72	0.00	0.0	145
	384.35	767.12	0.00		
14	100.10	73.74	0.00	0.0	145
	384.66	767.33	0.00		
15	99.97	73.79	121.81	133.0	145
	384.16	767.85	1539.95		
16	100.12	73.94	121.49	134.0	145
	384.74	769.41	1535.90		
17	99.99	73.60	121.49	n/r	n/r
	384.24	765.87	1535.90		
18	99.99	73.88	122.00	135.0	145
	384.24	768.78	1542.35		
19	100.04	73.45	121.74	0.0	145

	384.43	764.31	1539.06		
20	100.00	73.87	121.43	134.0	n/r
	384.28	768.68	1535.15		
21	99.95	73.45	121.12	132.0	145
	384.08	764.31	1531.23		

Table 10

Run #	Rxn Time (min)	Rxn Temp. (°C)
2	30	130-134
3	30	130-135
4	45	130-136
5	45	130-134
6	43	130-134
7	45	130-134
8	45	130-135
9	48	130-133
10	45	130-134
11	57	130-134
12	60	130-142
13	45	130-143
14	44	130-142
15	45	130-133
16	45	130-134
17	45	130-n/r
18	45	130-134
19	45	130-133
20	n/r	130 n/r
21	45	130 n/r

n/r indicates that no full temperature range or reaction time was recorded.

Example A6

Diacetone fructose (DAF) (780 kg; 780kg, Batch 1 and 2), pyridine (950.6 kg; 950.0 kg) and xylenes (1950 L; 1950 L) were mixed in a tank and the solution was transferred to a reactor with sulfamide (575 kg; 575 kg).

The mixture was heated to 128-133°C via stepwise heating. More specifically, the mixture was heated from 30 to 65°C in 80 minutes, then from 65 to 95°C in 60 minutes, followed by heating from 95 to 113°C in 60 minutes and 113 to 125°C in 85 minutes. The reactor jacket temperature was raised in 5°C increments to heat the batch to 128-133°C over 30 minutes and then held at the temperature for 30 minutes. The reaction mixture was then cooled at a rate of 30°C per hour to 40°C. To the mixture was then added purified water (1243L; 1240L). The mixture was cooled to 20°C before 50% NaOH (436 kg; 437 kg) solution was added. Organic and aqueous layers (product layers) were separated and the aqueous layer was filtered and transferred to another reactor. The organic layer was discarded. Purified water (1400L; 1400L) was added to the aqueous mixture and residual organics (pyridine and xylene) and water were distilled off under vacuum. The distillation was stopped after distilling a known volume (784 L; 784 L). Glacial acetic acid (214.5 kg; 224.4 kg) was added to neutralize the solution to pH 7. The mixture was seeded with topiramate seeds (1.0 kg; 1.0 kg) and was cooled to 0-5°C to crystallize the crude topiramate. The crude, wet topiramate (730 kg; 803.8 kg wet yield) was centrifuged and collected.

While the foregoing specification teaches the principles of the present invention, with examples provided

for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.